

Fermion dispersion in axion medium

N.V. Mikheev*, E.N. Narynskaya†

*Division of Theoretical Physics, Department of Physics,
Yaroslavl State University, Sovetskaya 14,
150000 Yaroslavl, Russian Federation.*

Abstract

The interaction of a fermion with the dense axion medium is investigated for the purpose of finding an axion medium effect on the fermion dispersion. It is shown that axion medium influence on the fermion dispersion under astrophysical conditions is negligible small if the correct Lagrangian of the axion-fermion interaction is used.

1 Introduction.

The investigations of elementary particle physics beyond the standard model are one of the interest line of modern researches. Particular attention is given to the light goldstone and pseudogoldstone bozons, arising at spontaneously breaking of some symmetry. In view of weak interaction with matter the special interest for such particles is caused by possible astrophysical applications. In particular, the nature of cool dark matter remains the mystery of modern cosmology. The one of the most likely candidate for the dark matter is axion associated with Peccei-Quinn symmetry [1,2]. The extremely small value of axion mass ¹,

$$10^{-6} \text{ eV} < m_a < 10^{-2} \text{ eV}, \quad (1)$$

and gigantic life time in vacuum [3]

$$\tau_{a \rightarrow 2\gamma} \sim 10^{50} \text{ sec} \left(\frac{F_a}{10^{10} \text{ GeV}} \right)^6 \left(\frac{\omega}{1 \text{ MeV}} \right), \quad (2)$$

are two paramount factors why axion is of astrophysical and cosmology interest. Axion with such mass (1) could form a considerable fraction of dark matter. Moreover the region with high density of axion (so-called axion stars) could exist in Universe. The possibility existence of such axion objects has been discussed [4]. Dense axion matter playing the role of active medium could influence on particles properties. In particular, the change of the dispersion relation of the high energy cosmic ray propagating in pseudoscalar particle medium was investigated in [5]. However, as it will be shown further the result presented in [5] is significantly overestimated. This is due to the fact that authors [5] used the Lagrangian with the Yukawa coupling as the Lagrangian of axion-fermion interaction.

*E-mail address:mikheev@uniyar.ac.ru

†E-mail address:elenan@uniyar.ac.ru

¹ We use natural units in which $c = \hbar = 1$.

In this letter we review the influence of cool axion matter on the fermion dispersion. The modification of the fermion dispersion is caused by fermion "forward-scattering" of medium axion. The axion-fermion interaction will be analyzed by using the Lagrangian in the form [6]

$$L_{af} = \frac{c_a}{F_a} \left(\bar{\psi} \gamma_\mu \gamma_5 \psi \right) \partial_\mu a, \quad (3)$$

where c_a is the model-dependent parameter of the order of unit, F_a is the Peccei-Quinn symmetry-breaking scale, ψ and a are the fermion and axion fields respectively. It should be noted that in order to obtain the correct amplitudes of processes involving two axions, it is necessary to use just the Lagrangian with derivative (3) rather than Yukawa Lagrangian

$$L_{af} = \frac{-2ic_a m_f}{F_a} \left(\bar{\psi} \gamma_5 \psi \right) a, \quad (4)$$

as it was indicated for the first time in [7].

2 Dispersion of fermion in axion medium.

For investigation the fermion dispersion in axion medium the process of fermion "forward scattering" of plasma axions (fig.1) should be taken into account. The amplitude of this process is intimately connected with fermion mass operator Σ by the following relation

$$M = -\bar{u}(p) \Sigma u(p), \quad (5)$$

where $u(p)$ is the bispinor amplitude of fermion.

The amplitude corresponding to the left diagram in fig.1 has the form

$$M^{(1)} = \frac{ic_a^2}{2F_a^2} \int \frac{d^3 q}{(2\pi)^3 \omega} f(\omega) \left(\bar{u}(p) [\hat{q} \gamma_5 S(p+q) \hat{q} \gamma_5] u(p) \right), \quad (6)$$

where $q^\mu = (\omega, \vec{q})$ and $p^\mu = (p_0, \vec{p})$ are the axion and fermion four-momentum correspondingly, $\hat{q} = (q_\mu \gamma^\mu)$, $f(\omega)$ is the axion distribution function, $f(\omega) = (e^{\omega/T} - 1)^{-1}$, $S(p)$ is the propagator of fermion with mass m_f :

$$S(p) = \frac{i(\hat{p} + m_f)}{p^2 - m_f^2}.$$




Figure 1: The diagrams describing fermion "forward scattering" of plasma axions.

The amplitude corresponding to the right diagram in fig.1 can be obtained from (6) by the substituting $q \rightarrow -q$. Performing some calculations, the total amplitude of the coherent scattering process $f + a \rightarrow f + a$ of all plasma axions can be written as

$$M = \frac{c_a^2}{2F_a^2} \int \frac{d^3q}{(2\pi)^3 \omega} f(\omega) \bar{u}(p) \left(\frac{(p^2 - m_f^2)\hat{q} + (\hat{p} + m_f)m_a^2}{(p+q)^2 - m_f^2} + q \rightarrow -q \right) u(p). \quad (7)$$

With the definition (5) the fermion mass operator has the form

$$\Sigma = \frac{-c_a^2}{2F_a^2} \int \frac{d^3q}{(2\pi)^3 \omega} f(\omega) \left(\frac{(p^2 - m_f^2)\hat{q} + (\hat{p} + m_f)m_a^2}{(p+q)^2 - m_f^2} + q \rightarrow -q \right). \quad (8)$$

There is good reason to believe that axion gas is most likely a cool nonrelativistic medium. So it is possible to carry out calculations in the approximation $q^\mu \simeq (m_a, \vec{q})$, $|\vec{q}| \ll m_a$. Under such conditions the expression for the mass operator is essentially simplified

$$\Sigma = \frac{-c_a^2 n_a}{4F_a^2 p_0^2 m_a} \left\{ 2p_0 \hat{U}(p^2 - m_f^2) - (\hat{p} + m_f)(p^2 - m_f^2 + m_a^2) \right\}, \quad (9)$$

where n_a is the concentration of axion gas, U^μ is the four-vector of medium which in its rest system has the form $U^\mu = (1, 0, 0, 0)$.

As one see from (9) the mass operator can be presented in the form:

$$\Sigma = Am_f + B\hat{p} + C\hat{U}, \quad (10)$$

$$A = B = \frac{n_a c_a^2 (p^2 - m_f^2 + m_a^2)}{4F_a^2 p_0^2 m_a}, \quad C = \frac{-n_a c_a^2 (p^2 - m_f^2)}{2F_a^2 p_0 m_a}.$$

The fermion dispersion law can be extracted from equation

$$\det(\hat{p} - m_f - \Sigma) = 0. \quad (11)$$

For the fermion energy in the cool axion medium from (11) with (10) one obtains

$$E_f \simeq \sqrt{\vec{p}^2 + m_f^2} + \frac{m_f^2(A+B)}{\sqrt{\vec{p}^2 + m_f^2}} - C, \quad (12)$$

where $\sqrt{\vec{p}^2 + m_f^2} = E_0$ is the fermion energy in vacuum.

This is our main result determining the fermion dispersion in the cool axion medium. For the purpose of comparison (12) with the previous result [5] we estimate the deviation of fermion mass squared in the cool axion medium from vacuum one, δm_f^2 :

$$\delta m_f^2 = p^2 - m_f^2 = E_f^2 - E_0^2 = E_f^2 - \vec{p}^2 - m_f^2.$$

The deviation of fermion mass squared from (12) can be written as

$$\delta m_f^2 \simeq 2m_f^2(A + B) - 2C\sqrt{\vec{p}^2 + m_f^2}. \quad (13)$$

Taking into account smallness of the parameters A, B and C the axion medium correction to squared of the fermion mass is

$$\delta m_f^2 \simeq \frac{n_a m_a c_a^2 m_f^2}{(\vec{p}^2 + m_f^2) F_a^2}. \quad (14)$$

It can be seen that this contribution is strongly suppressed not only by the energy scale of the breakdown symmetry F_a in denominator but the axion mass m_a in numerator as well. As was pointed above, the result for the squared of the fermion mass in axion medium under the same physical conditions was presented early [5] in the form

$$(\delta m_f^2)^{Sahu} \simeq \frac{n_a m_f^2}{2m_a F_a^2}. \quad (15)$$

The comparison of (14) with (15)

$$R = \frac{\delta m_f^2}{(\delta m_f^2)^{Sahu}} = \frac{2c_a^2 m_a^2}{E_0^2} \ll 1$$

illustrated that the result of the paper [5] is strongly overestimated. This is due to the fact authors [5] uncorrectly used the Lagrangian with pseudoscalar Yukawa coupling for investigation the process of fermion "forward scattering" of axions medium.

3 Conclusion

We have studied the influence of cool axion medium to the fermion dispersion. The contribution into squared of fermion mass due to presence of axion medium is calculated. The investigation of this type were performed early in paper [5]. In this paper we demonstrated that calculation of fermion dispersion in the axion medium with Yukawa Lagrangian leads to the significantly overestimate results. Under real astrophysical conditions the influence of axion medium to the fermion dispersion turns out to be negligible small.

This work supported in part by the Council on Grants by the President of Russian Federation for the Support of Young Russian Scientists and Leading Scientific Schools of Russian Federation under the Grant No. NSh-497.2008.2 and by the Russian Foundation for Basic Research under the Grant No. 07-02-00285-a.

References

- [1] R. D. Peccei, Quinn H.R., Phys.Rev.Lett. **38**, 1440 (1977); Phys.Rev. **D16**, 1791 (1977)
- [2] S. Weinberg, Phys.Rev.Lett. **40**, 223 (1978); F.Wilczek, Phys.Rev. **40**, 271 (1978)

- [3] G. G. Raffelt, Phys. Rep. **198**, 1 (1990).
- [4] C.J.Hogan and M.J.Rees, Phys.Rev.Lett. **B205**, 228 (1988); E.W.Kolb and I.I.Tkacher, Phys.Rev. **D49**, 5040 (1994).
- [5] S. Sahu, Mod.Phys.Lett.A **14**, 1287 (1999).
- [6] G.G. Raffelt, *Stars as Laboratories for Fundamental Physics* (University of Chicago Press, 1996).
- [7] G.G Raffelt, D. Seckel, Phys. Rev. Lett. **60**, 1793 (1988).